

A SLIGHT IMPROVEMENT TO GARAIEV'S SUM PRODUCT ESTIMATE

NETS HAWK KATZ AND CHUN-YEN SHEN

Indiana University

§0 INTRODUCTION

Let A be a subset of F_p , the field of p elements with p prime.

We let

$$A + A = \{a + b : a \in A, b \in A\},$$

and

$$AA = \{ab : a \in A, b \in A\}.$$

It is fun (and useful) to prove lower bounds on $\max(|A+A|, |AA|)$ (see e.g. [BKT],[BGK],[G]). Recently, Garaev [G] showed that when $|A| < p^{\frac{1}{2}}$ one has the estimate

$$\max(|A + A|, |AA|) \gtrsim |A|^{\frac{15}{14}}.$$

By using Plunneke's inequality in a slightly more sophisticated way, we improve this exponent to $\frac{14}{13}$. We believe that further improvements might be possible through aggressive use of Ruzsa covering.

§1 PRELIMINARIES

Throughout this paper A will denote a fixed set in the field F_p of p elements with p a prime. For B , any set, we will denote its cardinality by $|B|$.

Whenever X and Y are quantities we will use

$$X \lesssim Y,$$

1991 *Mathematics Subject Classification.* primary 42B25 secondary 60K35.

The first author was supported by NSF grant DMS 0432237.

to mean

$$X \leq CY,$$

where the constant C is universal (i.e. independent of p and A). The constant C may vary from line to line. We will use

$$X \lesssim Y,$$

to mean

$$X \leq C(\log |A|)^\alpha Y,$$

where C and α may vary from line to line but are universal.

We state some preliminary lemmas, mostly those stated by Garaev, but occasionally with different emphasis.

The first lemma is a consequence of the work of Glibichuk and Konyagin [GK]

Lemma 1.1. *Let $A_1 \subset F_p$ with $1 < |A_1| < p^{\frac{1}{2}}$. Then for any elements a_1, a_2, b_1, b_2 so that*

$$\frac{b_1 - b_2}{a_1 - a_2} + 1 \notin \frac{A_1 - A_1}{A_1 - A_1},$$

we have that for any $A' \subset A_1$ with $|A'| \gtrsim |A_1|$

$$|(a_1 - a_2)A' + (a_1 - a_2)A' + (b_1 - b_2)A'| \gtrsim |A_1|^2.$$

In particular such a_1, a_2, b_1, b_2 exist unless $\frac{A_1 - A_1}{A_1 - A_1} = F_p$. In case $\frac{A_1 - A_1}{A_1 - A_1} = F_p$, we may find $a_1, a_2, b_1, b_2 \in A_1$ so that

$$|(a_1 - a_2)A_1 + (b_1 - b_2)A_1| \gtrsim |A_1|^2.$$

Sketch of Proof. If $\frac{A_1 - A_1}{A_1 - A_1} \neq F_p$, it is immediate that there exist $a_1, a_2, b_1, b_2 \in A_1$ with $1 + \frac{b_1 - b_2}{a_1 - a_2} \notin \frac{A_1 - A_1}{A_1 - A_1}$. This automatically implies

$$|(a_1 - a_2)A' + (a_1 - a_2)A' + (b_1 - b_2)A'| \gtrsim |A_1|^2.$$

(See [GK]. If $x \notin \frac{A_1 - A_1}{A_1 - A_1}$ then each element of $A_1 + xA_1$ has but one representative $a + xa'$. On the other hand if

$$\frac{A - A}{A - A} = F_p,$$

then one can find $a_1, a_2, b_1, b_2 \in A_1$ so that $\frac{a_1 - a_2}{b_1 - b_2}$ has at most $|A|^2$ representatives as $\frac{a_3 - a_4}{b_3 - b_4}$ with $a_3, a_4, b_3, b_4 \in A$ which implies that $|A + \frac{a_1 - a_2}{b_1 - b_2}A|$ is large. Again, for more details see [GK]. $\square$

The following two lemmas, quoted by Garaev, are due to Ruzsa, may be found in [TV]. The first is usually referred to as Ruzsa's triangle inequality. The second is a form of Plunneke's inequality.

Lemma 1.2. *For any subsets X, Y, Z of F_p we have*

$$|Y - Z| \leq \frac{|Y - X||X - Z|}{|X|}.$$

Lemma 1.3. *Let $X, B_1, \dots, B_k$ be any subsets of F_p with*

$$|X + B_i| \leq \alpha_i |X|,$$

for i ranging from 1 to k . Then there exists $X_1 \subset X$ with

$$((1.1)) \quad |X_1 + B_1 + \dots + B_k| \leq \alpha_1 \dots \alpha_k |X_1|.$$

We record a number of Corollaries. The first two can be found in [TV]. The last one, we first became aware of in the paper of Garaev.

Corollary 1.4. *Let $X, B_1, \dots, B_k$ be any subsets of F_p . Then*

$$|B_1 + \dots + B_k| \leq \frac{|X + B_1| \dots |X + B_k|}{|X|^{k-1}}.$$

Proof. Simply bound $|B_1 + \dots + B_k|$ by $|X_1 + B_1 + \dots + B_k|$ and $|X_1|$ by $|X|$. $\square$

Corollary 1.4 is somewhat wasteful in that X_1 is unlikely to be both a singleton element and a set with the same cardinality as X . By applying Lemma 1.3 iteratively.

Corollary 1.5. *Let $X, B_1, \dots, B_k$ be any subsets of F_p . Then there is $X' \subset X$ with $|X'| > \frac{1}{2}|X|$ so that*

$$|X' + B_1 + \dots + B_k| \lesssim \frac{|X + B_1| \dots |X + B_k|}{|X|^{k-1}}.$$

Proof. Observe that for any $Y \subset X$ with $|Y| \geq \frac{|X|}{2}$, we have that

$$\frac{|Y + B_i|}{|Y|} \lesssim \frac{|X + B_i|}{|X|}.$$

Now recursively apply Lemma 1.3. That is, first apply it to $X, B_1, \dots, B_k$ obtaining a set X_1 satisfying

$$|X_1 + B_1 + \dots + B_k| \lesssim \frac{|X + B_1| \dots |X + B_k|}{|X|^k} |X_1|.$$

If $|X_1| > \frac{1}{2}|X|$ then stop and let $X' = X_1$. Otherwise apply Lemma 1.3 to $X \setminus X_1, B_1, \dots, B_k$. Proceeding recursively if $|X_1 \cup \dots \cup X_{j-1}| \geq \frac{1}{2}|X|$ then set

$$X' = X_1 \cup \dots \cup X_{j-1},$$

otherwise obtain the inequality

$$|X_j + B_1 + \dots + B_k| \lesssim \frac{|X + B_1| \dots |X + B_k|}{|X|^k} |X_j|.$$

Summing all the inequalities we obtained before stopping gives us the desired result. $\square$

Corollary 1.6. *Let $A \subset F_p$ and let $a, b \in A$. Then we have the inequalities*

$$|aA + bA| \leq \frac{|A + A|^2}{|aA \cap bA|},$$

and

$$|aA - bA| \leq \frac{|A + A|^2}{|aA \cap bA|}.$$

Proof. To get the first inequality, apply Corollary 1.4 with $k = 2$, $B_1 = aA$, $B_2 = bA$, and $X = aA \cap bA$.

To get the second inequality, apply Lemma 1.2 with $Y = aA$, $Z = -bA$ and $X = -(aA \cap bA)$. $\square$

§2 MODIFIED GARAEEV'S INEQUALITY

In this section, we slightly modify Garaev's argument to obtain

Theorem 2.1. *Let $A \subset F_p$ with $|A| < p^{\frac{1}{2}}$ then*

$$\max(|AA|, |A + A|) \gtrsim |A|^{\frac{14}{13}}.$$

Proof.

Following Garaev, we observe that

$$\sum_{a \in A} \sum_{b \in A} |aA \cap bA| \geq \frac{|A|^4}{|AA|}.$$

Therefore, we can find an element $b_0 \in A$, a subset $A_1 \subset A$ and a number N satisfying

$$|b_0A \cap aA| \approx N,$$

for every $a \in A_1$. Further

$$(2.1) \quad N \gtrsim \frac{|A|^2}{|AA|},$$

and

$$(2.2) \quad |A_1|N \gtrsim \frac{|A|^3}{|AA|}.$$

Now there are two cases. In the first case, we have

$$\frac{A_1 - A_1}{A_1 - A_1} = F_p.$$

If so, applying Lemma 1.1, we can find $a_1, a_2, b_1, b_2 \in A_1$ so that

$$|A_1|^2 \lesssim |(a_1 - a_2)A_1 + (b_1 - b_2)A_1| \leq |a_1A - a_2A + b_1A - b_2A|.$$

Applying Corollary 1.4 with $k = 4$ and with $B_1 = a_1A$, with $B_2 = -a_2A$ with $B_3 = b_1A$, with $B_4 = -b_2A$, and with $X = b_0A$. Then we apply Corollary 1.6 to bound above $|X + B_j|$. This yields

$$|A_1|^2 \lesssim \frac{|A + A|^8}{N^4|A|^3},$$

or

$$|A_1|^2 N^4 |A|^3 \lesssim |A + A|^8.$$

Applying (2.2), we get

$$(2.3) \quad N^2 |A|^9 \lesssim |A + A|^8 |AA|^2.$$

and applying (2.1), we get

$$(2.4) \quad |A|^{13} \lesssim |A + A|^8 |AA|^4.$$

The estimate (2.4) implies that

$$\max(|A + A|, |AA|) \gtrsim |A|^{\frac{13}{12}} \gtrsim |A|^{\frac{14}{13}},$$

so that we have more than we need in this case.

Thus we are left with the case that

$$\frac{A_1 - A_1}{A_1 - A_1} \neq F_p.$$

Thus we can find a_1, a_2, b_1, b_2 so that for any refinement $A' \subset A_1$ with $|A'| \gtrsim |A_1|$, we have

$$|A_1|^2 \lesssim |(a_1 - a_2)A' + (a_1 - a_2)A' + (b_1 - b_2)A'|.$$

Now we apply Corollary 1.5, choosing A' so that

$$|(a_1 - a_2)A' + (a_1 - a_2)A_1 + (b_1 - b_2)A_1| \lesssim \frac{|A + A| |(a_1 - a_2)A_1 + (b_1 - b_2)A_1|}{|A_1|}.$$

This is where we have improved over Garaev's original argument.

Then, as in the first case, estimating

$$|(a_1 - a_2)A_1 + (b_1 - b_2)A_1| \leq |a_1A - a_2A + b_1A - b_2A|,$$

and applying Corollary 1.4 with $X = b_0A$ and Corollary 1.6, we obtain

$$|A_1|^3 N^4 |A|^3 \lesssim |A + A|^9.$$

Applying (2.2), we get

$$(2.5) \quad N|A|^{12} \lesssim |A + A|^9 |AA|^3.$$

Now applying (2.1), we get

$$(2.6) \quad |A|^{14} \lesssim |A + A|^9 |AA|^4.$$

Inequality (2.6) proves the Theorem. $\square$

REFERENCES

- [BGK] Bourgain, J., Glibichuk, A.A., and Konyagin, S.V., *Estimates for the number of sums and products and for exponential sums in fields of prime order*, J. London Math. Soc. (2) **73** (2006), 380–398.
- [BKT] Bourgain, J., Katz, N, and Tao, T., *A sum product estimate in finite fields and Applications*, GAFA **14** (2004), 27–57.
- [G] Garaev, M.Z., *An explicit sum-product estimate in $\mathbb{F}_p$* , preprint.
- [GK] Glibichuk, A.A, and Konyagin, S.V, *Additive properties of product sets in fields of prime order*, preprint.
- [TV] Tao, T. and Vu, V., Cambridge Univ. Press (2006).